arXiv:1104.3617v1 [math.NT] 19 Apr 2011

Asymptotic harmonic behavior in the prime number distribution

Maurice H.P.M. van Putten

Le Studium IAS, 45071 Orléans Cedex 2, Université d'Orléans, France

ABSTRACT

Euler's identity is shown to give a relation between the zeros of the Riemann-zeta function and the prime numbers in terms of $\Phi(x) = x^{-\frac{1}{4}}\left[2\sqrt{x}\Sigma e^{-p^2\pi x}\ln(p) - 1\right]$ on $x > 0$, where the sum is over all primes p. If Φ is bounded on $x > 0$, then the Riemann hypothesis is true or there are infinitely many zeros Re $z_k > \frac{1}{2}$. The first 21 zeros give rise to asymptotic harmonic behavior in Φ defined by the prime numbers up to one trillion.

1. Introduction

The Riemann-zeta function $\xi(z)$ is a complex function of z defined in $\mathrm{Re}(z) > 1$ by the sum

$$\zeta(z) = 1 + \frac{1}{2^z} + \frac{1}{3^z} + \cdots \tag{1}$$

over the integers. The zeros z_k of $\zeta(z)$ comprise the negative even integers, $-2, -4, \cdots$, and an infinite distribution of nontrivial zeros $z_k = a_k + iy_k$ in the strip $0 < a_k < 1$.

A general approach to find zeros of as endpoints of continuation (Keller 1987). Here, we set out to analyze the z_k of as endpoints of continuation along a trajectory $z(\lambda)$, whose tangent $z'(\lambda)$ satisfies

$$\tau\frac{\zeta'(z)}{\zeta(z)} = -1 \tag{2}$$

in response to a choice of initial data $z(0) = z_0$, is a zero of $\xi(z)$. The trajectory $z(\lambda)$ continues whenever $\tau \neq 0$, i.e., whenever $\Theta(z) = -\frac{\zeta'(z)}{\zeta(z)}$ exists and is finite. Continuation halts only at a zero of $1/\Theta(z)$, corresponding to a zero of $\zeta(z)$, all of which are isolated. Continuation (2) can equivalently be formulated in terms of the prime numbers by

$$\xi(z) = \Sigma\ln(p)p^{-z} \quad (\mathrm{Re}\ z > 1), \tag{3}$$

representing the leading term the analytic extension of the sum

$$\Sigma\xi(mz) = \Sigma\frac{\ln(p)}{p^z - 1} = -\frac{\zeta'(z)}{\zeta(z)} \tag{4}$$

by virtue Euler's identity

$$\zeta(z) = \Pi\left(1 + p^{-z} + p^{-2z} + \cdots\right) = \Pi\left(1 - p^{-z}\right)^{-1}. \tag{5}$$

All known nontrivial zeros determined by numerical root finding satisfy Re $z_k = \frac{1}{2}$ to within numerical precision, the first three of which are $z_1 = \frac{1}{2} \pm 14.1347i$, $z_2 = \frac{1}{2} \pm 21.0220i$, $z_3 = \frac{1}{2} \pm 25.0109i$. By the symmetry

$$\zeta(s) = \zeta(1-s)\frac{\chi(1-s)}{\chi(s)}, \quad \chi(s) = \pi^{\frac{s}{2}}\Gamma\left(\frac{s}{2}\right), \quad \Gamma(z) = \int_0^\infty t^{z-1}e^{-t}dt, \tag{6}$$

it suffices to study zeros in the half plane $\mathrm{Re}(z) \geq \frac{1}{2}$. Fig. 1 illustrates root finding by (2) for the first few zeros.

In this paper, we study $\xi(z)$ in (3) in the relations between the distribution of primes and the location of z_k by Euler's identity in the expanded form

$$\xi(z) = -\frac{\zeta'(z)}{\zeta(z)} - \Sigma_{m\geq 2}\xi(mz). \tag{7}$$

The results will be formulated in terms of a density and normalized density function of the primes defined on the positive real numbers $x > 0$, given by

$$\Phi(x) = \frac{2\sqrt{x}\phi(x) - 1}{x^{\frac{1}{4}}}, \quad \phi(x) = \Sigma e^{-p^2\pi x}\log(p), \tag{8}$$

where summation in $\phi(x)$ is over all primes. The function $\xi(z)$ has the property that it possesses poles at the zeros z_k of $\zeta(z)$. In particular, the pole at $z = 1$ is a consequence of the Prime Number Theorem applied to the Chebyshev function (Dusart 1999), by which $2\sqrt{x}\phi(x) - 1 = o(1)$ as x approaches zero. In our application of a Fourier transform to (7), the z_k introduce a sum of residues

$$Z(\lambda) = \Sigma\gamma_k e^{-\lambda(z_k - \frac{1}{2})} \tag{9}$$

defined for all $\lambda < 0$, where the

$$\gamma_k = \gamma(z_k), \quad \gamma(z) = \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}}, \tag{10}$$

are absolutely summable by Stirling's formula and the asymptotic distribution of z_k.



Fig. 1.— Shown are the trajectories of continuation $z(\lambda)$ in the complex plane z by numerical integration of (2) with initial data $z_0 = 1 + ni$ ($n = 1, 2, 3, \cdots$) indicated by small dots on Re(z)=1. Continuation produces roots indicated by open circles, defined by finite endpoints of $z(\lambda)$ in the limit as λ approaches infinity. The roots produced by the choice of initial data are the first three on Re $z = \frac{1}{2}$ and -2 and -4 of the trivial roots.

Theorem 1.1. *In the limit as $x > 0$ becomes small, we have the asymptotic behavior*

$$\Phi(x) + \left[\frac{\Gamma\left(\frac{1}{4}\right)}{2\pi^{\frac{1}{4}}} + Z\left(\ln\sqrt{x}\right)\right] + \frac{\Gamma\left(\frac{1}{6}\right)}{3\pi^{\frac{1}{6}}}x^{\frac{1}{12}} = o\left(x^{\frac{1}{12}}\right). \tag{11}$$

Corollary 1.2. *If Φ is bounded, then the Riemann hypothesis is true or there are infinitely many zeros Re $z_k > \frac{1}{2}$.*

A similar relation between the distribution of z_k and the primes is given by (Hadamard 1893; von Mangoldt 1895)

$$\frac{\psi(u) - u}{\sqrt{u}} = -\Sigma\frac{u^{z_k - \frac{1}{2}}}{z_k} - \frac{\ln(2\pi) + \ln\sqrt{1 - u^{-2}}}{\sqrt{u}}, \tag{12}$$

based on the Chebyshev functions

$$\psi(u) = \Sigma_{p^k \le u}\ln(p), \quad \vartheta(u) = \Sigma_{p \le u}\ln p, \tag{13}$$

where the sum is over all primes p and integers k. In Theorem 1.1, normalization of $\Phi(x)$ in Theorem 1.1 is by $x^{\frac{1}{4}}$ according to (8) and Z is absolutely convergent for all $x > 0$, whereas in (12) normalization is by $\sqrt{u}$ and the sum $\Sigma \frac{u^{z_k - \frac{1}{2}}}{z_k}$ is not absolutely convergent. Similar to Corollary 1.2, it will be appreciated that the left hand side of (12) will be bounded in the limit of large u if the Riemann hypothesis is true.

In §2, we give some background on $\zeta(z)$. In §3 we introduce an integral representation of $\xi(z)$ and derive one of its integral properties associated with the singularity of $\zeta(z)$ at $z = 1$. In §4, we apply Cauchy's integral formula to $\zeta(z)$ to derive a sum of residues associated by the z_k. The proof Theorem 1.1 is given in §5 by a Fourier transform and asymptotic analysis of the expanded Euler's identity. In §6, we report on a direct evaluation of $\Phi(x)$ using the primes up to one trillion, to show the harmonic behavior as as encoded in Z by the first few zeros z_k. We summarize our findings in §7.

2. Background

Our analysis uses some well known properties of $\zeta(z)$ and its zeros $z_k = a_k + iy_k$, e.g., in Titchmarsh (1986); Lehmer (1988); Dusart (1999); Keiper (1992); Ford (2002)).

Riemann provided an analytic extension of $\xi(z)$ to the entire complex plane by expressing each term n^{-z} in terms of $\Gamma\left(\frac{z}{2}\right)$,

$$\zeta(z) = \frac{\pi^{\frac{z}{2}}}{\Gamma\left(\frac{z}{2}\right)} \int_0^\infty x^{\frac{z}{2}-1}\theta_1(x)dx, \tag{14}$$

where

$$\theta_1(x) = \frac{\theta(x) - 1}{2}, \quad \theta(x) = \Sigma_{-\infty}^{\infty} e^{-n^2\pi x} \tag{15}$$

satisfies the asymptotic property $\theta_1(x) \sim \frac{1}{2\sqrt{x}}$ as x approaches zero by the identity $\theta(x^{-1}) = \sqrt{x}\theta(x)$ for the Jacobi function $\theta(x)$.[1] Evaluation of the integral (14) on Re $z > 1$ hereby reveals the meromorphic expression (e.g. Borwein et al. 2006)

$$\zeta(z) = \frac{\pi^{\frac{z}{2}}}{\Gamma\left(\frac{z}{2}\right)} \left\{ \frac{1}{z(z-1)} + f(z) \right\}, \quad f(z) = \int_1^\infty \left(x^{\frac{z}{2}-1} + x^{-\frac{z}{2}-\frac{1}{2}} \right) \theta_1(x)dx, \tag{16}$$

---

[1] When $z = n$ is an integer, the pre-factor $\frac{\pi^{\frac{n}{2}}}{\Gamma\left(\frac{n}{2}\right)}$ will be recognized to be half the surface area $S(n)$ of a unit $n-$sphere.

representing a maximal analytic continuation of $\zeta(z)$, and the presence of a simple pole at $z=1$ with residue 1.

Riemann also introduced the symmetric form $Q(z)\zeta(z)$, $Q(z) = \frac{1}{2}z(z-1)\pi^{-\frac{z}{2}}\Gamma\left(\frac{z}{2}\right)$ satisfying $Q(z)\zeta(z) = Q(1-z)\zeta(1-z)$, whereby

$$\zeta(z) = \pi^{z-1}\zeta(1-z)\frac{\Gamma(\frac{1}{2}-\frac{z}{2})}{\Gamma(\frac{z}{2})} = \frac{\pi^z\zeta(1-z)}{\cos\left(\frac{1}{2}\pi z\right)\Gamma(\frac{z}{2})\Gamma(\frac{1}{2}+\frac{z}{2})} = \frac{\pi^{z-\frac{1}{2}}2^{z-1}\zeta(1-z)}{\cos\left(\frac{1}{2}\pi z\right)\Gamma(z)} \tag{17}$$

using $\Gamma(\frac{1}{2}-\frac{z}{2})\Gamma(\frac{1}{2}+\frac{z}{2}) = \frac{\pi}{\cos(\pi z)}$ and $\Gamma(z)\Gamma(z+\frac{1}{2}) = 2^{1-2z}\sqrt{\pi}\Gamma(2z)$. Along the vertical line $z = 1-iy$, $\zeta(z) \neq 0$ (Littlewood 1922, 1924, 1927; Wintner 1941), which allows consideration of its logarithmic derivative,

$$\frac{\zeta'(z)}{\zeta(z)} = -\frac{\zeta'(1-z)}{\zeta(1-z)} - \ln(2\pi) + \frac{\pi}{2}\tan\left(\frac{\pi z}{2}\right) - \psi(z) \tag{18}$$

in terms of the digamma function

$$\psi(z) = \frac{\Gamma'(z)}{\Gamma(z)} \sim \ln(z) + O(z^{-1}) \tag{19}$$

in the limit of large $|z|$.

Lemma 2.1. *In the limit of large y, the logarithmic derivative of $\zeta(z)$ satisfies*

$$\frac{\zeta'(iy)}{\zeta(iy)} = \frac{\zeta'(1-iy)}{\zeta(1-iy)} + O(\ln y). \tag{20}$$

Proof. The result follows from (19) and (18). □

Lemma 2.2. *Along the line $z = iy$, we have the asymptotic expansion $|\gamma(iy)| \sim \sqrt{\frac{2\pi}{y}}e^{-\frac{\pi}{2}y}$ in the limit of large y, whereby the γ_k are absolutely summable.*

Proof. Recall (10) and the asymptotic expansion $\Gamma(z) = \sqrt{2\pi}z^{z-\frac{1}{2}}e^{-z}\left[1+O(z^{-1})\right]$ with a branch cut along the negative real axis. In the limit of large y_k, and the asymptotic behavior of the zeros z_k, whereby $y_k \sim \frac{2\pi k}{\ln k}$, and hence $|\gamma_k| \sim e^{-\frac{\pi^2 k}{\ln k}}$,where we used $|\arg z_k| \sim \frac{\pi}{2}$ in the limit of large k. It follows that the γ_k are absolutely summable. Numerically, their sum is rather small, $\Sigma|\gamma_k| = 3.5\times10^{-5}$ based on a large number of known zeros z_k. □

Lemma 2.3. *In the limit of large y, we have*

$$\left|\gamma(iy)\frac{\zeta'(iy)}{\zeta(iy)}\right| = O\left(y^{-\frac{1}{4}}e^{-\frac{\pi}{2}y}\ln y\right). \tag{21}$$

Proof. By Lemma 2.1-2, we have

$$\left|\gamma(iy)\frac{\zeta'(iy)}{\zeta(iy)}\right| \leq |\gamma(iy)| \left(\left|\frac{\zeta'(1-iy)}{\zeta(1-iy)}\right| + |A(y)| \right) \sim \sqrt{\frac{2\pi}{y}} \left(\left|\frac{\zeta'(1-iy)}{\zeta(1-iy)}\right| + O(\ln y) \right) e^{-\frac{\pi}{2}y} \quad (22)$$

asymptotically for large y. By (Richert 1967; Cheng 1999; Titchmarsh 1986; Broughan 2005)

$$\left|\frac{\zeta'(1-iy)}{\zeta(1-iy)}\right| \leq c(\ln y)^{\frac{2}{3}}(\ln\ln y)^{\frac{1}{3}} \quad (23)$$

on $y > \delta$ for some positive constants c, δ. □

These observations give rise to the following.

Proposition 2.4. *The Fourier transform*

$$L(\lambda) = PV \int_{-\infty}^{\infty} \gamma(iy)\frac{\zeta'(iy)}{\zeta(iy)} e^{-i\lambda y}\frac{dy}{2\pi} + \ln(2\pi), \quad (24)$$

where PV refers to the Principle Value about $y = 0$*, is well defined and* $F(\lambda) \to 0$ *in the limit of large* λ*.*

Proof. The result follows from Lemma 2.1-3 and the Riemann-Lebesque lemma. □

3. An integral representation of $\xi(z)$

We begin with an integral representation of $\xi(z)$ in terms of $\phi(x)$ and $\Phi(x)$ given in (8), following steps very similar to those by Riemann leading to (16). To this end, we define $g(z) = g_1(z) + g_2(z)$,

$$g_1(z) = \int_{-\infty}^{0} \Phi(e^{2\lambda})e^{\lambda(z-\frac{1}{2})}d\lambda, \quad g_2(z) = 2\int_{0}^{\infty} \phi(e^{2\lambda})e^{\lambda z}d\lambda \quad (25)$$

for Re $z > 1$ and, respectively, all z.

Proposition 3.1. *On* Re $z > 1$*, we have*

$$\Sigma \ln(p)p^{-z} = \frac{\pi^{\frac{z}{2}}}{\Gamma\left(\frac{z}{2}\right)} \left\{ \frac{1}{z-1} + g(z) \right\}. \quad (26)$$

Proof. Following steps similar to those leading to (16), we have $\xi(z) = \frac{\pi^{\frac{z}{2}}}{\Gamma(\frac{z}{2})}\int_0^\infty x^{\frac{z}{2}-1}\phi(x)dx$, that may be decomposed in integration over $(0, 1]$ and $[1, \infty)$,

$$g_1(z) = \frac{1}{2}\int_0^1 x^{\frac{2z-1}{4}}\Phi(x)\frac{dx}{x}, \quad g_2(z) = \int_1^\infty x^{\frac{z}{2}-1}\phi(x)dx, \quad (27)$$

where $g_1(z)$ includes regularization about $z = 1$ made explicit in (26) by $\frac{1}{z-1}$, where the singularity at $z = 1$ in ξ can be seen to result from the Prime Number Theorem in the form of $2\sqrt{x}\phi(x) - 1 = o(1)$ on the basis of the asymptotic behavior of the function (Dusart 1999). The substitution $x = e^{2\lambda}$ gives (25). □.

Lemma 3.2. *The analytic extension of* $g_1(z)$ *extends to the real values* $z > \frac{1}{2}$.

Proof. In a neighborhood of $0 < z < 1$, we may write

$$-\frac{\zeta'(z)}{\zeta(z)} = \frac{\pi^{\frac{z}{2}}}{\Gamma\left(\frac{z}{2}\right)}\left[\frac{1}{z-1} + u_1(z)\right], \tag{28}$$

where $u_1(z)$ is analytic about $z > 0$. With $z = a + ib$, the second term on the right hand side in the expanded Euler's identity (7) satisfies

$$\Sigma_{m\geq 3}|\xi(mz)| \leq \Sigma_{n\geq 3}\frac{\log(n)n^{-3a}}{1-n^{-a}} < -\frac{\sqrt{2}\zeta'(3a)}{\sqrt{2}-1}, \tag{29}$$

whereby it is bounded in Re $z = a > \frac{1}{2}$. Since the second term $\zeta(2z)$ in (7) is analytic in Re $z = a > \frac{1}{2}$, it follows that $g(a)$ as defined in Proposition 3.1 is analytic on $a > \frac{1}{2}$. In view of the analytic and finite behavior of the right hand side in (7), the first and second term on the left hand side in (7) remain balanced as a approaches $\frac{1}{2}$ from the right, giving

$$\xi(a) = -\frac{1}{2a-1} + u_2(a), \tag{30}$$

where $u_2(a)$ is analytic at $a = \frac{1}{2}$. As a approaches $\frac{1}{2}$ from the right, we have

$$g_1(a) = \frac{1}{2a-1} + u_3(a), \tag{31}$$

where $u_3(a)$ is analytic about $a = \frac{1}{2}$. □

By Proposition 3.2, the behavior of the sign of $\Phi(x)$ in a neighborhood of $x = 0$ is relevant. If $\Phi(x)$ is of one sign in some neighborhood, then $g_1(z)$ has an analytic extension into Re $z > \frac{1}{2}$ with no singularities, implying the absence of z_k here. However, this requires information on the point wise behavior of $\Phi(x)$, which goes beyond the relatively weaker integrability property (31). To make a step in this direction, we next apply a linear transform to (7) to derive the asymptotic behavior of $\Phi(x)$ in terms of the distribution z_k.

4. A sum of residues Z associated with the non-trivial zeros

We next define

$$h(z) = \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}}\frac{\zeta'(z)}{\zeta(z)} + \frac{1}{z-1} \tag{32}$$

and its Fourier transform

$$H(\lambda) = \int_{a-i\infty}^{a+i\infty} h(z) e^{-\lambda z} \frac{dz}{2\pi i}. \tag{33}$$

Lemma 4.1. *The function $h(z)$ has a simple pole at $z = 1$ with residue 1 and simple poles at each of the nontrivial zeros z_k of $\zeta(z)$ with residue γ_k.*

Proof. As a consequence of (e.g. Borwein et al. 2006)

$$\zeta_1(z) = \frac{1}{2} z(z-1) \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}} \zeta(z), \quad \frac{\zeta_1'(z)}{\zeta_1(z)} = B + \Sigma_k \left(\frac{1}{z - z_k} + \frac{1}{z_k} \right), \tag{34}$$

where B is a constant, so that

$$\frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}} \frac{\zeta'(z)}{\zeta(z)} + \frac{1}{z-1} = \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}} \left[B + \Sigma_k \left(\frac{1}{z - z_k} + \frac{1}{z_k} \right) \right] + A. \tag{35}$$

Here

$$A = \frac{1}{z-1} \left(1 - \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}} \right) - \frac{\Gamma\left(\frac{z}{2}\right)}{z \pi^{\frac{z}{2}}} - \frac{2\Gamma'\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}} \ln \pi} \tag{36}$$

includes contributions from the logarithmic derivative of the factor to $\zeta(z)$ in (34), whose singularities are restricted to the trivial zeros of $\zeta(z)$. □

Proposition 4.2. *The Fourier transform of $h(z)$ over Re $z > \sup a_k$ satisfies*

$$H(\lambda) = e^{-\frac{\lambda}{2}} Z(\lambda) + O(1) \tag{37}$$

in the limit of large $\lambda < 0$.

Proof. Consider contour integration over the rectangle Re $z = 0$, Re $z = a$ and $z = x \pm iY$ $(0 < x < a)$. The line integral over $z = 0$ satisfies $O\left(e^{\frac{\lambda}{2}}\right)$ by Proposition 2.4. Integration over $z = x + iY$ $(0 < x < a)$ gives

$$e^{-iY} \int_{iY}^{iY+a} h(z) e^{-\lambda x} \frac{dx}{2\pi i} = e^{-iY} \int_{iY}^{iY+a} \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}} \frac{\zeta'(z)}{\zeta(z)} \frac{dx}{2\pi i} + O\left(Y^{-1}\right), \tag{38}$$

where we choose Y to be between two consecutive values of y_k. We have

$$\int_{iY}^{iY+a} \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}} \frac{\zeta'(z)}{\zeta(z)} \frac{dx}{2\pi i} \sim \gamma_k \int_{iY}^{iY+a} \frac{\zeta'(z)}{\zeta(z)} \frac{dx}{2\pi i} \sim \frac{\gamma_k}{2\pi i} \ln(2a-1) \left[1 + \frac{4ai(y_k - Y)}{1 - 2a} + \pi i \right]. \tag{39}$$

In the limit as k approaches infinity, $y_k - Y$ approaches zero and $|\gamma_k|$ becomes small by Lemma 2.2., whence

$$\left(\int_{iY}^{iY+a} - \int_{-iY}^{-iY+a}\right) \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}} \frac{\zeta'(z)}{\zeta(z)} \frac{dx}{2\pi i} \sim \ln(2a-1)\text{Im }\gamma_k = O\left(\ln(2a-1)\sqrt{\frac{2\pi}{y_k}}\, e^{-\frac{\pi}{2}y_k}\right). \tag{40}$$

The result now follows in the limit as k approaches infinity, taking into account the residue sum $e^{-\frac{\lambda}{2}}Z(\lambda)$ associated with the z_k and absolute summability of the γ_k. $\square$

5. Proof of Theorem 1.1

Following the expanded Euler identity (7), we define

$$f(z) = -g_2(z) - h(z) - \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}}\Sigma_{m\geq 2}\xi(mz) \tag{41}$$

for all z away from the z_k, its Fourier transform

$$F(\lambda) = \int_{a-i\infty}^{a+i\infty} f(z)e^{-\lambda z}\frac{dz}{2\pi i} \tag{42}$$

on $a > \sup a_k$, and the coefficients

$$c_m(z) = \frac{\pi^{\frac{mz}{2}}\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}\Gamma\left(\frac{mz}{2}\right)}, \quad C_m = m^{-1}\pi^{-\frac{1}{2m}}\Gamma\left(\frac{1}{2m}\right). \tag{43}$$

for all $m \geq 1$. We remark that in particular, $C_2 = \frac{\Gamma(\frac{1}{4})}{2\pi^{\frac{1}{4}}}$ and $m^{-1}c_m(m^{-1}z) = 1 + \left(\frac{1}{2}\ln\pi + \frac{1}{2}\gamma\right)z + O\left(z^2\right)$ has a well defined limit and $C_m \to 2$ in the limit as m becomes arbitrarily large.

Lemma 5.1. *The decomposition of the sum* $\Sigma_{m\geq 1}\xi(mz) = \xi(z) + \Sigma_{m\geq 2}\xi(mz)$ *is well-defined on Re* $z > \frac{1}{2}$.

Proof. By the Prime Number Theorem, $p_k \sim k\ln(k)$, whereby summation over the tails $k \geq n$ defined by large n satisfy

$$\Sigma_{k\geq n}^{\infty}\frac{\ln(p_k)}{p_k^{2a}} \sim \Sigma_{k\geq n}^{\infty}\left[\frac{1}{k^{2a}\ln(k)^{2a-1}} + \frac{\ln\ln(k)}{(k\ln(k))^{2a}}\right] < \infty \tag{44}$$

whenever $a > \frac{1}{2}$. Hence, for $z = a + iy$, $|\Sigma\ln(p)p^{-2z}| \leq \Sigma\ln(p)p^{-2a} < \infty$ whenever $a > \frac{1}{2}$. It follows that $|\Sigma_{m\geq 2}\xi(mz)| \leq \Sigma_{m\geq 2}\Sigma_p p^{-(m-2)a}\ln(p)p^{-2a} \leq \Sigma_{m\geq 0}2^{-m}\Sigma_p\ln(p)p^{-2a} = \Sigma_p\ln(p)p^{-2a} < \infty$ on Re $z > \frac{1}{2}$, whereby the sum $\Sigma_{m\geq 2}\xi(mz)$ is well-defined. Given the analytic continuation of $\Sigma_{m\geq 1}\xi(mz)$ to Re $z > \frac{1}{2}$ by the logarithmic derivative of the (analytic continuation of) $\zeta(z)$, it follows that the (analytic continuation of) $\xi(z)$ is well-defined. □

Lemma 5.2. $F(\lambda) = e^{-\frac{\lambda}{2}}\Phi(e^{2\lambda})$.

Proof. By Proposition 3.1, the Fourier transform

$$\int_{a-\infty}^{a+i\infty} g_1(z)e^{-\lambda z}\frac{dz}{2\pi i} \tag{45}$$

exists uniquely on Re $z > 1$. Here it equals $F(\lambda)$, since $f(z)$ is the analytic extension in Re $z > \frac{1}{2}$ of $g_1(z)$ in Re $z > 1$. □

Lemma 5.3. *For any $m \geq 2$, the Fourier transform of $\frac{c_m(z)}{mz-1}$ over Re $z = a > \frac{1}{2}$ satisfies*

$$D_m(\lambda) = C_m e^{-\frac{\lambda}{m}} + o(1) \tag{46}$$

Proof. We have

$$D_m(\lambda) = e^{\frac{\lambda}{2}}\int_{-\infty}^{\infty}\frac{c_m(iy)}{imy-1}e^{-i\lambda y}\frac{dy}{2\pi} + C_m e^{\lambda\left(\frac{1}{2}-\frac{1}{m}\right)}. \tag{47}$$

The integral (47) exists by virtue of a removable singularity of $c_m(z)$ at $z = 0$ and decays along the imaginary axis when $m \geq 2$ by the Riemann-Lebesque lemma. □

We now consider the Fourier transform of the expanded Euler identity in the form (41) over Re $z = a$ on a term by term basis. To organize terms, we define

$$r_N(z) = g_2(z) + \Sigma_{m\geq 2}c_m(z)g(mz) + \frac{\Gamma\left(\frac{z}{2}\right)}{\pi^{\frac{z}{2}}}\Sigma_{m\geq N+1}\xi(mz), \tag{48}$$

whereby

$$f(z) + \Sigma_{m=2}^{N}\frac{c_m(z)}{mz-1} + r_N(z) = -h(z), \tag{49}$$

and the Fourier transform

$$R_N(\lambda) = \int_{a-i\infty}^{a+i\infty} r_N(z)e^{-\lambda z}\frac{dz}{2\pi i}. \tag{50}$$

Lemma 5.4. $R_N(\lambda) = o(1)$ *in the limit of large $\lambda < 0$.*

Proof. Since $r_N(z)$ is analytic in Re $z > \frac{1}{2}$, its transform $R_N(\lambda)$ exists and is uniquely defined for $a > \frac{1}{2}$. The result follows from the Riemann-Lebesque lemma. □

Proof of Theorem 1.1. The Fourier transform of the expanded Euler identity in the form (49) is

$$F(\lambda) + D_2(\lambda) + D_3(\lambda) + R_3(\lambda) + H(\lambda) = 0. \tag{51}$$

By Propositions 5.1, 4.2 and Lemmas 5.1-5.4, we have

$$e^{-\frac{\lambda}{2}}\Phi(e^{2\lambda}) + C_2 e^{-\frac{\lambda}{2}} + C_3 e^{-\frac{\lambda}{3}} + e^{-\frac{\lambda}{2}} Z(\lambda) = O(1) \tag{52}$$

With $x = e^{2\lambda}$, Theorem 1.1 now follows. □

6. Numerical illustration using the primes up to one trillion

Fig. 2 shows the first harmonic in the normalized density $\Phi(x)$ for small x evaluated for the 37.6 billion primes up to $y = 10^{12}$, allowing x down to 2.6×10^{-23} ($\lambda = -26$) in view of the requirement for an accurate truncation in $\phi(x)$ as defined by (8). The harmonic behavior is emergent in

$$R(x) = \Phi(x) - C_2 - C_3 x^{\frac{1}{12}}. \tag{53}$$

To search for higher harmonics $Z_i(\lambda)$ associated with the zeros z_i in $\lambda\epsilon[-26, -11.7759] = [\lambda_1 - \lambda_2, \lambda_1 + \lambda_2]$, we compare the spectrum of $\Phi\left(e^{2\lambda}\right)$ by taking a Fast Fourier Transform with respect to α,

$$\lambda(\alpha) = \lambda_1 + \lambda_2 \cos\alpha \ \ (\alpha\epsilon[0, 2\pi]), \tag{54}$$

and compare the results with an analytic expression for the Fourier coefficients of the $Z_i(\lambda)$ $(i = 1, 2, \cdots)$,

$$c_{ni}[\lambda_1, \lambda_2] = 2\mathrm{Re}\{(-i)^n \gamma_i e^{-i\lambda_1 z_i} J_n\left(-\lambda_2 z_i\right)\}, \tag{55}$$

where $J_n(z)$ denotes the Bessel function of the first of order n. Fig. 3 shows the first 21 harmonics in our evaluation of $\Phi(x)$, which appears to be about the maximum that can be calculated by direct summation in quad precision.

7. Conclusions

The zeros $z_k = a_k + iy_k$ of the Riemann-zeta function are endpoints of continuation: $\xi(z_k) = 0$ iff $1/\Theta(z_k) = 0$. By Euler's identity (5), continuation and hence the endpoints are controlled by an integral $\xi(z)$ of a regularized sum $\Phi(e^{2\lambda})$ over the prime numbers as defined by (8).

The contributions of the zeros z_k introduce asymptotic harmonic behavior in $\Phi\left(e^{2\lambda}\right)$ as a function of $\lambda < 0$ as defined by the sum $Z(\lambda)$ of residues associated with the z_k, shown in Figs. 1-2. This harmonic behavior in the prime number distribution appears to be exceptionally dense: primes up to 4 billion are needed to identify the first 4 harmonics, up to 70 billion for the 10 and up to 1 trillion for the first 21. It appears that the prime number range scales exponentially with the number of harmonics it contains. Conversely, knowledge of a relatively small number of zeros z_k provides information on exponentially large primes numbers.

Theorem 1.1 introduces a correlation between Φ, defined over the primes, and Z, defined over the z_k. Suppose there are a finite number of zeros z_k in Re $z > \frac{1}{2}$. We may then consider k^* for which $a_{k^*} = \max a_k$ gives rise to dominant exponential growth in $Z(\lambda)$ in the limit as $\lambda < 0$ becomes large. This observation leads to Corollary 1.2. Z can remain bounded in $x > 0$ only if the Riemann hypothesis is true, or if $Z(\lambda)$ remains fortuitously bounded as an infinite sum over $a_k > \frac{1}{2}$ with no maximum in $a < 1$.

Acknowledgment. The author gratefully acknowledges stimulating discussions with Anton F.P. van Putten and insightful comments from Fabian Ziltener. Some of the manuscript was prepared at the Korea Institute for Advanced Study, Dongdaemun-Gu, Seoul. This research was supported in part by the National Science Foundation through TeraGrid resources provided by Purdue University under grant number TG-DMS100033. We specifically acknowledge the assistance of Vicki Halberstadt, Rich Raymond and Kimberly Dillman. The computations have been carried out using Lahey Fortran 95.

REFERENCES

Borwein, P., Choi, S., Rooney, B., & Weirathmueller, A., 2006, *The Riemann Hypothesis* (Berlin: Springer Verlag)

Broughan, K.A., 2005, preprint

Cheng, Y., 1999, Rocky Mountain J. Math. 29, 115

Dusart, P.. 1999, C. R. Math. Rep. Acad. Sci. Canad 21, 53

Ford, K., 2002, in Number Theory for the Millenium (Urbana, IL, 2000), 2, 25

Hadamard, J., J. Math. Pures Appl., 171

Keiper, J.B., Math. Comput., 58, 765

Keller, H.B., 1987, *Numerical Methods in Bifurcation Problems* (Berlin: Springer Verlag/Tata Institute for Fundamental Research)

Lehmer, D.H., 1988, Math. Comput., 50, 265

Littlewood, J.E., 1922, Proc. London Math. Soc., ser. 2, 20, xxii

Littlewood, J.E., 1924, Proc. London Math. Soc., ser. 2, 24, 175

Littlewood, J.E., 1927, Proc. London Math. Soc., ser. 2, 27, 349

Richert, H.E., 1967, Math. Ann. 169, 97

von Mangoldt, H., J. Reine Angew. Math. 114, 255

Titchmarsh, E.C., 1986, The theory of the Riemann Zeta-function, Oxford, 2nd ed. (revised by D. R. Heath-Brown)

Wintner, A., 1941, Amer. J. Math., 63, 575

This preprint was prepared with the AAS LaTeX macros v5.0.



Fig. 2.— The top window shows the normalized density $\Phi\left(e^{2\lambda}\right)$ on $\lambda\epsilon[-26,-11.7756]$ and its leading order approximation $1.3616+1.5332e^{\frac{\lambda}{6}}$. The asymptotic harmonic behavior is apparent in the residual difference (53) between the two, shown in the bottom two windows, including the period of 2.2496 in λ associated with the first zero $z_*=\frac{1}{2}\pm 14.1347i$.



Fig. 3.— Shown are the absolute values of the Fourier coefficients $c_n[\lambda_1, \lambda_2]$ of $\Phi\left(e^{2\lambda}\right)$ obtained by a Fast Fourier Transform (FFT) of (53) on the computational domain (54), where $\lambda_1 = -26$, $\lambda_2 = -11.7756]$ covers 32 periods of $Z_1(\lambda)$ (*dots*), on the basis of the 37,607,912,2019 primes up to 1,000,000,000,0039. The resulting spectrum is compared with the exact spectra $c_{ni}[\lambda_1, \lambda_2]$ of the $Z_i(\lambda)$ given by the analytic expression (55) for $i = 1, 2, 3, \cdots$ (*continuous line*). Shown are also the individual spectra of $Z_i(\lambda)$ for $i = 1, 8$ and 15 associated with the zeros z_1, z_8 and z_{15}. The match between the computed and exact spectra accurately identifies the first 21 harmonics of $Z(\lambda)$ in Φ out of 22 shown, corresponding to the first 21 nontrivial zeros z_i of $\zeta(z)$.